



Articles

Perceived Value of Virtual Reality Simulations to Gain Insights into Delirium: A Pilot Study

Laura Fontanesi¹, Arzu Çöltekin¹, Thomas Kunz², Sandro Arrigoni¹, Steven Bourke¹, Alasdair MacLulich¹, Rasita Vinay³, Tobias Kowatsch^{3,4}, Nico Julian Zobrist³, Clara Julia Lampe⁵, Martina Schöpf⁶, Jeremy Howick⁷, Mathias Schlögl⁶

¹ FHNW University of Applied Sciences and Arts, ² Draeger Schweiz AG, ³ University of Zurich, ⁴ University of St.Gallen, ⁵ Technical University of Munich, ⁶ Clinic Barmelweid, ⁷ University of Leicester

Keywords: Delirium, Virtual Reality, Immersive Simulation, Empathy Training, Medical Education, Nursing Education

<https://doi.org/10.56392/001c.158141>

Delirium

Background

Delirium affects one-in-four hospitalized individuals aged 65-years or older, and is associated with serious complications, including post-traumatic stress disorder, prolonged hospitalization, and increased mortality. Healthcare professionals often lack insight into the distress that patients experience; this is known to undermine empathic care and timely detection. Virtual reality (VR) is emerging as a promising tool to enhance empathy and understanding of the patient experience due to its inherent immersive, embodied, and experience-based characteristics.

Objective

To pilot and obtain preliminary subjective evidence on the usability, empathic impact, and educational value of a VR simulation grounded in qualitative evidence of patients' delirium experiences.

Methods

23 university students completed a 7-minute VR delirium simulation and an immediate 42-item survey. This included usability items adapted from the System Usability Scale, items conceptually informed by different validated questionnaires, and study-specific items. Full instruments were not administered to minimize respondent burden and target context-specific elements. Spearman correlations explored response patterns. Open-ended items were summarized descriptively.

Results

Twelve of 22 Likert items differed from neutrality (3) after correction. Participants found the simulation intuitive ($M=4.30$, $SD=0.70$, $d=1.86$) and its immersive nature engaging ($M=4.22$, $SD=0.74$, $d=1.65$), reporting increased empathy ($M=4.39$, $SD=0.58$, $d=2.39$), improved understanding of delirium-related emotional distress ($M=4.13$, $SD=0.81$, $d=1.48$) and cognitive distortions ($M=4.22$, $SD=0.60$, $d=1.81$). They endorsed the integration of VR into health education and expressed interest in potential additional features. Qualitative responses highlighted VR's value for empathy training and to improve doctor-patient communication, while calling for even greater realism and personalization.

Conclusion

This pilot study suggests that VR simulations are perceived as effective, engaging tools for enhancing empathy and insight in delirium-related education and care training.

INTRODUCTION

Delirium is a common and severe neuropsychiatric syndrome characterized by an acute onset of cognitive disturbances that cannot be attributed to a pre-existing or evolving neurocognitive disorder.¹⁻³ Approximately one in four hospitalized adults over 65 experience delirium.⁴ It is as-

sociated with increased mortality, prolonged hospital stay, long-term cognitive decline, institutionalization, and psychological distress, including post-traumatic stress disorder, for both patients and caregivers.^{1,2,5}

Despite the availability of validated screening tools, delirium is frequently unrecognized or misdiagnosed, particularly in its hypoactive form.³ From the patient perspec-

tive, episodes may be difficult to articulate, as acknowledging confusion or “losing one’s mind” during hospitalization adds to an already vulnerable situation. Qualitative research describes fear, shame, and difficulties in communication, further complicating diagnoses.⁶ Nurses, who often conduct initial assessments, report barriers such as time constraints, inadequate training, and limited awareness of baseline cognition. Knowledge gaps are not restricted to nursing staff: medical students also receive inconsistent delirium education and often underestimate its consequences.⁷

Traditional delirium education mostly consists of face-to-face education, and, to a lesser extent, e-learning and interprofessional education.⁸ To facilitate the translation from theory to practice, there is growing interest in experiential methods such as virtual reality (VR) simulations,⁵ which immerse learners in the perspective of patients undergoing cognitive or sensory disruptions.^{9–11} VR has been shown to facilitate perspective-taking and increase empathy.^{12–14} Large-scale studies suggest VR simulations lead to more durable behavioral changes than traditional approaches,¹³ and meta-analytic evidence indicates that VR is particularly effective at enhancing perspective-taking, though effects on empathy are more variable.¹⁴ This variability may depend on how VR is implemented, as VR is generally regarded as a highly effective tool for fostering empathy.^{12,15,16}

In clinical education, VR simulations have been used to foster empathic communication and understanding of patient needs.^{17–21} For example, VR simulations of dementia or sensory loss improved students’ comprehension of cognitive impairment and appreciation of patient perspectives.²² However, delirium has received little attention in VR-based education. A small study with nursing students found VR delirium simulations useful,²¹ and a quasi-experimental study with 18 nurses showed improvements in empathy and knowledge compared to lecture-based training.²³ Yet these studies were limited by small samples and scenarios not explicitly grounded in extended accounts of patient experiences.

With this study, we contribute to the emerging delirium VR research area by evaluating the perceived impact of VR delirium simulations comprising a few key scenes, grounded in qualitative evidence of patients’ delirium experiences²⁴ (which emphasized disorientation, fear, helplessness, and altered perception as main experiences) and input from clinical experts. Through post-simulation survey data, we explored how participants interpreted the simulation’s educational relevance, empathic impact, and potential value for healthcare training.

METHODS

PARTICIPANTS AND PROCEDURE

We recruited 23 university students attending an educational session organized with the Swiss Study Foundation, bringing together university students from diverse disciplinary backgrounds interested in delirium care (as future

caregivers) personally or professionally, to deepen their understanding of delirium and to evaluate our immersive VR simulation. Inclusion criteria were age ≥ 18 , German fluency, and no known contraindications to VR (contraindications to VR included a history of epilepsy or severe motion sickness).

Sessions were conducted in a lecture hall configured for parallel VR use, with eleven Meta Quest 3 headsets set up at individual stations to prevent interference. After screening and providing informed consent, participants received a brief orientation, completed a 7-minute VR simulation, and then filled out a 42-item questionnaire on their own devices.

ETHICAL STATEMENT

The study was reviewed using a checklist, according to institutional procedures at the FHNW School of Applied Psychology, and was classified as not requiring cantonal ethics approval (in Switzerland, human subjects research is overseen by cantonal ethics committees). The study adhered to GDPR requirements.

VR SIMULATION

The simulation was developed in Unity and deployed on Meta Quest 3 headsets with integrated audio. Informed by qualitative syntheses of patient experiences²⁴ and clinical expert input, the simulation followed a linear, three-scene hospital narrative (**Figure 1**): introduction by a nurse; delirium phase with restraint in bed and visual/auditory distortions (e.g., distorted lights, a threatening zombie-like figure, a crawling spider, staff speaking as if the patient were absent); and recovery phase with improved lighting and empathetic staff behavior, underscoring the impact of caregiver interactions.

MEASUREMENTS

Participants completed a study-specific 42-item questionnaire, covering five domains: usability/immersion, cognitive and clinical learning, emotional engagement/empathy, ethical/public-health considerations, and future directions, plus demographics. Six items adapted the wording of the System Usability Scale²⁵ (SUS); because the full SUS was not administered, no SUS score was computed. While this approach did not allow us to infer validated constructs, it prioritized context relevance and reduced participants’ burden.²⁶ Additional items were conceptually inspired by the User Engagement Scale²⁷ (UES), Immersive Tendencies Questionnaire²⁸ (ITQ), and Toronto Empathy Questionnaire²⁹ (TEQ), but were tailored to VR-based delirium training; accordingly, no UES/ITQ/TEQ scores were calculated. Most items (22) used five-point Likert-type responses, and 7 open-ended questions invited further elaboration after each topic. The open-ended questions were inspired by direct clinical observations as well as the prior work. The full questionnaire can be seen as a Supplementary Material.




Figure 1. Key scenes from the VR simulation of delirium.

Screenshots illustrate core elements of the VR experience. Top-left: the participant's perspective while lying in a dimly lit hospital room, restrained at the wrist. Top-right: a zombie-like figure enters the room, representing visual hallucinations. Bottom-left: a doctor speaks while the room fills with fog, simulating perceptual disorientation and emotional detachment. The script was drafted and refined by delirium specialists to reflect typical clinical situations and communication patterns before being implemented in the simulation, and speech was simulated using an off-the-shelf speech engine. Bottom-right: a spider crawls on the participant's virtual arm during an injection, evoking tactile hallucinations and fear.

STATISTICAL ANALYSES

The Likert items were summarized descriptively. To test departures from the neutral midpoint (3), we ran two-tailed one-sample t-tests with Bonferroni correction across the 22 inferential items ($\alpha_{\text{per-test}}=0.05/22$). While we treated Likert-type responses as approximately continuous for inference,³⁰ we reported Shapiro-Wilk and Wilcoxon signed-rank tests as robustness checks and provided confidence intervals and effect sizes.³¹ To explore response patterns, we computed Spearman inter-item correlations and visualized a correlation matrix; given the adapted nature of our questionnaire, these analyses are exploratory and not intended to validate original scales.

RESULTS

PARTICIPANT CHARACTERISTICS

The sample consisted of 23 university students (74% aged 18-24; 26% aged 25-34; 56% women). 43% percent reported a medical background; while most did not have professional healthcare experience (52%), 35% had professional healthcare experience and 13% had experience as informal caregivers. Prior exposure to delirium was 48% none, 39% direct, 13% indirect. Regarding VR, 65% were occasional users (21% first-time; 13% frequent). None had previously experienced a delirium-focused VR simulation. [Table 1](#) details the qualifiers used to describe the participants.

QUANTITATIVE RESULTS (LIKERT ITEMS)

All participants who agreed to join the experiment completed the tasks, i.e., the response rate was 100%, and overall ratings were positive ([Table 2](#)). Because Shapiro-Wilk

Table 1. Participant characteristics (was presented as Section 1 in the questionnaire)

Participant characteristic	Related question(s)
Medical background	What is your academic or professional background? <input type="checkbox"/> Medicine <input type="checkbox"/> Psychology <input type="checkbox"/> Computer Science <input type="checkbox"/> Public Health <input type="checkbox"/> Engineering <input type="checkbox"/> Social Sciences <input type="checkbox"/> Other (please specify): _____
Professional healthcare experience / informal caregiver	Do you have any professional experience in healthcare and/or patient care? <input type="checkbox"/> Yes, as a medical professional <input type="checkbox"/> Yes, as a caregiver <input type="checkbox"/> Yes, both <input type="checkbox"/> No
Prior exposure to delirium	Have you ever had personal or professional experience with delirium (as a patient, caregiver, or clinician)? <input type="checkbox"/> Yes <input type="checkbox"/> I'm familiar but only indirectly <input type="checkbox"/> No
VR experience	Have you had prior experience with Virtual Reality? <input type="checkbox"/> Yes, frequently <input type="checkbox"/> Occasionally <input type="checkbox"/> No, this will be my first time
Experience with delirium-focused VR simulation	Have you had prior experience of delirium scenes in Virtual Reality? <input type="checkbox"/> Yes, frequently <input type="checkbox"/> Occasionally <input type="checkbox"/> No, this will be my first time If yes, how has your previous exposure to delirium influenced your experience with the VR simulation? (Open-ended) Response: _____

indicated non-normality for most items, **Wilcoxon signed-rank tests** were reported alongside **one-sample t-tests**, though that did not change inferences for any item.

Usability and immersion. Participants found the simulation 'intuitive and easy to navigate' ($M=4.30$, $SD=0.70$, $p<.0023$), its immersive nature engaging ($M=4.22$, $SD=0.74$, $p<.0023$), and the sensory effects 'effective in simulating a delirium experience' ($M=4.00$, $SD=0.67$, $p<.0023$). Items implying negative experiences were near neutrality: about the simulation being 'cumbersome' ($M=2.30$, $SD=1.29$, $p=.017$), 'taxing' ($M=2.65$, $SD=1.07$, $p=.134$), needing 'technical support' ($M=2.22$, $SD=1.13$, $p=.003$), and finding 'too much inconsistency in the sensory effects' ($M=2.61$, $SD=0.94$, $p=.059$).

Cognitive and clinical learning. Participants agreed that the VR simulation 'provided valuable insights into cognitive distortions' ($M=4.22$, $SD=0.60$, $p<.0023$) and didn't find it 'unnecessarily complex' ($M=1.65$, $SD=0.57$, $p<.0023$). Responses to an improved 'ability to identify signs of delirium experience' were above the midpoint but not significant ($M=3.61$, $SD=0.94$, $p=.005$), while responses to an improved 'ability to differentiate between delirium and dementia' were neutral ($M=3.09$, $SD=1.02$, $p=.680$).

Emotional engagement and empathy. Participants reported an increase in 'empathy toward delirium patients' ($M=4.39$, $SD=0.58$, $p<.0023$) and improved understanding of 'the emotional distress of delirium patients' ($M=4.13$, $SD=0.81$, $p<.0023$). They disagreed with 'viewing the experience of delirium patients in VR did not touch me' ($M=2.00$, $SD=1.09$, $p<.0023$). Feeling 'emotionally overwhelmed or anxious during the simulation' was near neutrality ($M=2.61$, $SD=1.12$, $p=.107$), as well as 'viewing the experience was too synthetic (game-like) to create empathy' ($M=3.00$, $SD=1.04$, $p=1.0$).

Ethical and public health considerations. Participants agreed that the simulation 'should be incorporated into medical and caregiver training programs' ($M=4.57$, $SD=0.51$, $p<.0023$) and 'could help reduce disparities in delirium care and education' ($M=4.39$, $SD=0.50$, $p<.0023$). 'There are ethical concerns regarding the emotional intensity of VR-based delirium simulations' was below the midpoint, but not sig-

nificant ($M=2.30$, $SD=1.02$, $p=.003$). 'I have concerns about the effort / costs involved in building such a system vs. its benefits' was neutral ($M=2.57$, $SD=1.08$, $p=.066$).

Future directions for research. Participants agreed with 'future VR studies should explore how delirium symptoms evolve over time' ($M=4.13$, $SD=0.63$, $p<.0023$) and 'VR-based multi-user simulations could improve interdisciplinary teamwork' ($M=4.48$, $SD=0.67$, $p<.0023$).

Inter-item correlations. Exploratory Spearman correlations (Figure 2) showed a few patterns in participant responses. Participants who reported an empathy increase also tended to report higher understanding of delirium: The reported increase in empathy correlated both with the understanding delirium patients' emotional distress ($\rho=.57$) and cognitive distortions ($\rho=.48$); understanding delirium patients' emotional distress and their cognitive distortions also correlated with each other ($\rho=.35$). Reporting increased empathy was also positively correlated with the belief that the simulation could reduce disparities in care ($\rho=.47$) and should be used in training ($\rho=.34$), and was negatively correlated with 'viewing the experience of delirium patients in VR did not touch me' ($\rho=-.51$).

In contrast, participants who found the simulation less user-friendly, rating it as 'cumbersome', 'taxing', or 'unnecessarily complex', did not report higher understanding: Finding the VR simulation 'unnecessarily complex' negatively correlated with the understanding of delirium patients' emotional distress ($\rho=-.42$). Finding it 'cumbersome' was correlated with finding it 'taxing' ($\rho=.49$) and finding the simulations' sensory effects effective ($\rho=.42$), suggesting some appreciation of sensory realism despite cognitive load. However, these usability concerns did not correlate strongly ($|\rho| \geq .40$) with endorsement (using it in training, its potential to reduce disparities and multi-user applications).

QUALITATIVE RESULTS (OPEN-ENDED ITEMS)

21 participants responded to the question about whether 'AI and adaptive/personalized VR could improve the realism and effectiveness of this training'. Many mentioned that per-

Table 2. Item-level descriptive statistics, confidence intervals, and inferential tests for the quantitative items of the study questionnaire

Section	Item (as in the questionnaire)	Mean	SD	95% CI (low)	95% CI (high)	p-value (t-test)	p-value (Wilcoxon)	Cohen's d	Significance
Section 2: Usability and VR experience	The VR simulation was intuitive and easy to navigate	4.30	0.70	4.00	4.61	0.0000	0.0001	1.86	Yes (Above 3)
	I found VR simulation cumbersome to use	2.30	1.29	1.74	2.86	0.0172	0.0263	-0.54	No
	The immersive nature of the simulation contributed to my engagement	4.22	0.74	3.90	4.54	0.0000	0.0001	1.65	Yes (Above 3)
	Using this VR simulation was taxing	2.65	1.07	2.19	3.12	0.1335	0.1341	-0.33	No
	The sensory effects (visual distortions, auditory cues) used in the VR were effective in simulating a delirium experience	4.00	0.67	3.71	4.29	0.0000	0.0001	1.48	Yes (Above 3)
	I thought there was too much inconsistency in the sensory effects	2.61	0.94	2.20	3.02	0.0586	0.0634	-0.42	No
	I think that I would need the support of a technical person to be able to use this VR simulation	2.22	1.13	1.73	2.70	0.0030	0.0048	-0.70	No
Section 3: Cognitive and Clinical Impact	I believe the VR experience improved my ability to identify signs of delirium experience: After seeing these scenes, I believe I can more easily tell if a patient is experiencing delirium	3.61	0.94	3.20	4.02	0.0052	0.0093	0.65	No
	The VR simulation provided valuable insights into cognitive distortions experienced by delirium patients	4.22	0.60	3.96	4.48	0.0000	0.0000	2.03	Yes (Above 3)
	I found the VR simulation unnecessarily complex	1.65	0.57	1.40	1.90	0.0000	0.0000	-2.35	Yes (Below 3)
Section 4: Emotional and Empathy-Based Impact	My ability to differentiate between delirium and dementia improved after the VR experience	3.09	1.02	2.64	3.54	0.6800	0.6736	0.09	No
	The VR simulation increased my empathy toward delirium patients	4.39	0.58	4.14	4.64	0.0000	0.0000	2.39	Yes (Above 3)
	I felt emotionally overwhelmed or anxious during the simulation	2.61	1.12	2.13	3.09	0.1073	0.1058	-0.35	No
	Viewing the experience of delirium patients in VR did not touch me	2.00	1.09	1.53	2.47	0.0002	0.0015	-0.92	Yes (Below 3)
	Viewing the experience was too synthetic (game-like) to create empathy	3.00	1.04	2.55	3.45	1.0000	1.0000	0.00	No
	The VR experience helped me understand the emotional distress of delirium patients more effectively than conventional exchanges (e.g., direct storytelling)	4.13	0.81	3.78	4.48	0.0000	0.0001	1.39	Yes (Above 3)
Section 5: Public Health and Ethical Considerations	I think the VR simulation should be incorporated into medical and caregiver training programs	4.57	0.51	4.35	4.78	0.0000	0.0000	3.09	Yes (Above 3)
	There are ethical concerns regarding the emotional intensity of VR-based delirium simulations	2.30	1.02	1.86	2.75	0.0035	0.0058	-0.68	No
	This VR approach could help reduce disparities in delirium care and education	4.39	0.50	4.18	4.61	0.0000	0.0000	2.79	Yes (Above 3)
Section 6: Future Research	I have concerns about the effort / costs involved in building such a system vs. its benefits	2.57	1.08	2.10	3.03	0.0665	0.0717	-0.40	No
	Future VR studies should explore how delirium symptoms evolve over time	4.13	0.63	3.86	4.40	0.0000	0.0000	1.81	Yes (Above 3)
	VR-based multi-user simulations could improve interdisciplinary teamwork	4.48	0.67	4.19	4.77	0.0000	0.0000	2.22	Yes (Above 3)

Statistics for each survey item are shown: mean, standard deviation (SD), and 95% confidence interval (CI) of the mean. One-sample t-tests (two-tailed) compare means against the neutral midpoint of 3; Wilcoxon signed-rank tests are included as robustness checks. Cohen's d effect size expresses the magnitude of the difference of the mean from the neutral point. Bonferroni-adjusted significance threshold for the t-tests is $\alpha = 0.05/22$. Items are reported with the exact survey wording.




Figure 2. Spearman correlations among the the quantitative items of the study questionnaire

Heatmap of Spearman correlation coefficients between individual questionnaire items. Item labels were shortened for readability in the figure and reflect the main content of each question (e.g., “Intuitive” = “The VR simulation was intuitive and easy to navigate”); they are not intended as validated constructs. Correlation coefficients are presented without significance testing. Colors represent direction and magnitude (red = positive, blue = negative).

sonalization could increase realism and make the simulation more immersive (“By interacting with individual patients based on their own needs, fears and experiences, one could certainly enhance the experience. An AI algorithm could explore these personal traits.”). Enhancements in sensory immersion and interactivity were also recommended. 22 participants responded to the question about whether this simulation could be used for ‘early delirium diagnosis in clinical settings’. The most frequent response was that awareness could help diagnoses, but that the simulation is not a diagnostic tool per se (“To raise awareness of delirium and make people consider it as a differential diagnosis, help people to have more empathy for delirious patients”). 20 participants responded to the question about whether the simulation could improve ‘doctor-patient communication and bedside manner’. Most highlighted that VR could increase empathy by helping clinicians ‘see how patients might react’ and ‘have a better idea of what patients experience’. All 23 participants responded to the question about whether they would ‘recommend this simulation for patient or caregiver education’. The majority said they would recommend the simulation, often noting that it was simple and effective for caregivers to better understand patients’ confusion (‘because an experience is more memorable than a lecture’) and that it would benefit from more realism.

Finally, 19 participants gave suggestions on future research directions. Ideas included adding the interaction with family members, multi-sensorial experiences, and more scenarios and patient-doctors interactions. Several comments suggested developing more interactive and realistic scenarios adapted to different patients’ feedback. 18

participants responded to the question about new features or improvements for future VR simulations. Frequent suggestions were increasing realism (e.g., graphics, movement, smell), making the simulation less game-like, and expanding interaction possibilities.

DISCUSSION

This pilot study adds to a growing body of research supporting the use of VR to foster empathy and clinical insight in health education. Participants reported increased empathy and greater insight into patients’ cognitive and emotional disturbances, consistent with prior work on VR-supported perspective-taking and engagement in clinical training.^{12, 13} The correlation between reported empathy increase and improved understanding aligns with evidence that affect can support learning.^{32,33} Lasting learning outcomes can be achieved through the use of stories (referred to as *narrative medicine*³⁴), which arguably is most beneficial to underserved communities, e.g., those with low literacy or cognitive impairment. Furthermore, if the realism levels are carefully crafted, there is evidence in prior work in other contexts that virtual environments can offer memorable experiences optimized for cognitive load concerns.^{35,36} Confirming the need for such optimization, in our pilot study, usability mattered: While usability ratings were overall positive, participants who perceived higher complexity or burden tended to report lower empathy and insight, highlighting the need to design emotionally powerful but technically accessible VR experiences. Qualitative responses emphasized the value of simulating the patient perspective

and recommended further personalization, increased sensory realism, and inclusion of varied delirium presentations. Taken together, these findings from our pilot study provide early evidence that a concise, qualitatively grounded VR simulation of delirium is both feasible and valued by learners (not just experienced practitioners^{5,23}), offering a strong foundation for scaled, patient-centred training. Some limitations need to be acknowledged. We used a study-specific, adapted questionnaire rather than validated scales; the small convenience sample of students, shared testing space, and reliance on immediate self-report limit generalizability. Notably, while reported empathy and insight increased, differentiation between delirium and dementia remained near neutral, suggesting that empathy-oriented VR may need to be complemented with targeted diagnostic training.

Future work following this pilot study should enroll clinically active learners, compare VR with active pedagogical controls, and incorporate behavioral outcomes (e.g., screening frequency/accuracy, recognition time, and communication quality). Testing adaptive (e.g., personalized depth of field simulation and motion control against cybersickness), interactive, and multi-user/team-based designs, as well as automated approaches to VR content creation to improve scalability, and comparative studies to alternative versions e.g., smartphone-based augmented reality (AR) versions are also envisioned. With these refinements, VR can complement conventional education by conveying the subjective experience of delirium while supporting practice-relevant learning.

ACKNOWLEDGEMENTS

We are grateful for our colleagues and students who joined us in the data collection efforts; and the event hosts, Dr. Sandra Bastelberger, Clara Lampe, and Nico Zobrist, for facilitating participant recruitment, as well as the use of the venue by the University of Zurich. We are thankful to the Swiss Study Foundation (Dr. Sandra Bastelberger) for having organized the educational event.

AUTHOR CONTRIBUTIONS

Laura Fontanesi: Investigation; Formal analysis; Writing – original draft; Writing – review & editing; Visualization.

Arzu Çöltekin: Conceptualization; Methodology; Resources; Writing – review & editing; Supervision; Project administration; Funding acquisition.

Thomas Kunz: Conceptualization; Software; Investigation; Resources; Writing – review & editing.

Sandro Arrigoni: Software; Investigation.

Mathias Schlögl: Conceptualization; Resources; Writing – review & editing; Supervision; Funding acquisition.

Steven Bourke: Writing – review & editing.

Alasdair MacLulich: Writing – review & editing.

Rasita Vinay: Writing – review & editing.

Tobias Kowtasch: Writing – review & editing.

Nico Zobrist: Writing – review & editing.

Clara Lampe: Writing – review & editing.

Martina Schöpf: Writing – review & editing.

Vincenza Frisardi: Writing – review & editing.

FUNDING SOURCES

The study was partially funded by Swiss Innovation Agency Innosuisse (project number: 39337.1 IP-ICT) for the members of the FHNW, as well as Draeger Schweiz AG and Clinic Barmelweid resources.

DECLARATION OF INTERESTS

RV and TK are affiliated with the Centre for Digital Health Interventions (CDHI), a joint initiative of the Institute for Implementation Science in Health Care, University of Zurich, the Department of Management, Technology, and Economics at ETH Zurich, and the Institute of Technology Management and School of Medicine at the University of St Gallen. CDHI is funded in part by the Swiss health insurer CSS, the Austrian health care provider (and corporate start-up of UNIQA) Mavie Next, and the Swiss investor MTIP. TK was also a co-founder of Pathmate Technologies, a university spin-off company that creates and delivers digital clinical pathways. However, neither CSS, Mavie Next, Pathmate Technologies nor MTIP were involved in this study. Furthermore, TK has neither shares of Pathmate Technologies nor any formal role in the company. NJZ receives a scholarship from the Swiss Study Foundation, funded by the Werner Siemens Foundation for the academic year 25/26. SB is the owner and founder of PersonalPulse. SB is a fellow of EUPATI foundation. CJL receives a scholarship from the German Academic Scholarship Foundation. All other authors declare no competing interests.

Accepted: January 28, 2026 CEST. Published: May 04, 2026 CEST.



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-4.0). View this license's legal deed at <http://creativecommons.org/licenses/by/4.0> and legal code at <http://creativecommons.org/licenses/by/4.0/legalcode> for more information.

REFERENCES

1. Inouye SK, Westendorp RGJ, Saczynski JS. Delirium in elderly people. *Lancet*. 2014;383(9920):911-922. doi:[10.1016/S0140-6736\(13\)60688-1](https://doi.org/10.1016/S0140-6736(13)60688-1)
2. Fong TG, Tulebaev SR, Inouye SK. Delirium in elderly adults: diagnosis, prevention and treatment. *Nat Rev Neurol*. 2009;5(4):210-220. doi:[10.1038/nrneuro.2009.24](https://doi.org/10.1038/nrneuro.2009.24)
3. Martins S, Fernandes L. Delirium in Elderly People: A Review. *Front Neurol*. 2012;3. doi:[10.3389/fneur.2012.00101](https://doi.org/10.3389/fneur.2012.00101)
4. Gibb K, Seeley A, Quinn T, et al. The consistent burden in published estimates of delirium occurrence in medical inpatients over four decades: a systematic review and meta-analysis study. *Age Ageing*. 2020;49(3):352-360. doi:[10.1093/ageing/afaa040](https://doi.org/10.1093/ageing/afaa040)
5. Schlögl M, Fontanesi L, Çöltekin A, et al. Immersive Virtual Reality Simulation to Promote Empathy in Delirium Care: A Pilot Evaluation. *Delirium*. Published online November 24, 2025. doi:[10.56392/001c.154095](https://doi.org/10.56392/001c.154095)
6. Van Rompaey B, Van Hoof A, van Bogaert P, Timmermans O, Dilles T. The patient's perception of a delirium: A qualitative research in a Belgian intensive care unit. *Intensive Crit Care Nurs*. 2016;32:66-74. doi:[10.1016/j.iccn.2015.03.002](https://doi.org/10.1016/j.iccn.2015.03.002)
7. Rosenstein B. IS DELIRIUM EDUCATION IMPORTANT: MEDICAL STUDENTS' ATTITUDES FOLLOWING A DELIRIUM WORKSHOP. *Innov Aging*. 2024;8(Suppl 1):1042. doi:[10.1093/geroni/igae098.3353](https://doi.org/10.1093/geroni/igae098.3353)
8. Lee SY, Fisher J, Wand APF, et al. Developing delirium best practice: a systematic review of education interventions for healthcare professionals working in inpatient settings. *Eur Geriatr Med*. 2020;11(1):1-32. doi:[10.1007/s41999-019-00278-x](https://doi.org/10.1007/s41999-019-00278-x)
9. Pottle J. Virtual reality and the transformation of medical education. *Future Healthc J*. 2019;6(3):181-185. doi:[10.7861/fhj.2019-0036](https://doi.org/10.7861/fhj.2019-0036)
10. Jiang H, Vimalasvaran S, Wang JK, Lim KB, Mogali SR, Car LT. Virtual Reality in Medical Students' Education: Scoping Review. *JMIR Med Educ*. 2022;8(1):e34860. doi:[10.2196/34860](https://doi.org/10.2196/34860)
11. Hirt J, Beer T. Use and impact of virtual reality simulation in dementia care education: A scoping review. *Nurse Educ Today*. 2020;84:104207. doi:[10.1016/j.nedt.2019.104207](https://doi.org/10.1016/j.nedt.2019.104207)
12. Barbot B, Kaufman JC. What makes immersive virtual reality the ultimate empathy machine? Discerning the underlying mechanisms of change. *Comput Hum Behav*. 2020;111:106431. doi:[10.1016/j.chb.2020.106431](https://doi.org/10.1016/j.chb.2020.106431)
13. Herrera F, Bailenson J, Weisz E, Ogle E, Zaki J. Building long-term empathy: A large-scale comparison of traditional and virtual reality perspective-taking. *PLOS ONE*. 2018;13(10):e0204494. doi:[10.1371/journal.pone.0204494](https://doi.org/10.1371/journal.pone.0204494)
14. Ventura S, Badenes-Ribera L, Herrero R, Cebolla A, Galiana L, Baños R. Virtual Reality as a Medium to Elicit Empathy: A Meta-Analysis. *Cyberpsychology Behav Soc Netw*. 2020;23(10):667-676. doi:[10.1089/cyber.2019.0681](https://doi.org/10.1089/cyber.2019.0681)
15. Hassan R. DIGITALITY, VIRTUAL REALITY AND THE "EMPATHY MACHINE." *Digit Journal*. 2020;8(2):195-212. doi:[10.1080/21670811.2018.1517604](https://doi.org/10.1080/21670811.2018.1517604)
16. Bujić M, Salminen M, Macey J, Hamari J. "Empathy machine": how virtual reality affects human rights attitudes. *Internet Res*. 2020;30(5):1407-1425. doi:[10.1108/INTR-07-2019-0306](https://doi.org/10.1108/INTR-07-2019-0306)
17. Bard JT, Chung HK, Shaia JK, Wellman LL, Elzie CA. Increased medical student understanding of dementia through virtual embodiment. *Gerontol Geriatr Educ*. 2023;44(2):211-222. doi:[10.1080/02701960.2022.2067850](https://doi.org/10.1080/02701960.2022.2067850)
18. Villmore DL, Dyer EJ, Gugliucci MR. Teaching empathy: comparison of a virtual reality experience using head-mounted display versus group streaming. *Gerontol Geriatr Educ*. 2024;0(0):1-12. doi:[10.1080/02701960.2024.2328514](https://doi.org/10.1080/02701960.2024.2328514)
19. Dyer E, Swartzlander BJ, Gugliucci MR. Using virtual reality in medical education to teach empathy. *J Med Libr Assoc JMLA*. 2018;106(4):498-500. doi:[10.5195/jmla.2018.518](https://doi.org/10.5195/jmla.2018.518)
20. Tuohy D, Boland P, Stark P, et al. Digital education about delirium for health care professional students: a mixed methods systematic review. *BMC Med Educ*. 2024;24:762. doi:[10.1186/s12909-024-05725-3](https://doi.org/10.1186/s12909-024-05725-3)

21. Matsuura J, Kunii T, Teranishi K, Tatsuana H, Noborio H. Construction of a Postoperative Delirium Experience System for Nursing Student Education Using VR and HMD. *IEEJ Trans Electron Inf Syst.* 2022;142(5):536-542. doi:[10.1541/ieejieiss.142.536](https://doi.org/10.1541/ieejieiss.142.536)
22. VR Training Library | Immersive Training | Embodied Labs. Accessed May 17, 2025. <https://www.embodiedlabs.com/immersive-experiences-library-overview>
23. Günther T, Schimböck F, Grosch M, Nydahl P. Virtual reality for delirium: immersive training courses promote empathy and expertise in delirium management among registered nurses. *Intensive Crit Care Nurs.* 2025;91:104167. doi:[10.1016/j.iccn.2025.104167](https://doi.org/10.1016/j.iccn.2025.104167)
24. Kuusisto-Gussmann E, Höckelmann C, Von Der Lühe V, Schmädig R, Baltés M, Stephan A. Patients' experiences of delirium: A systematic review and meta-summary of qualitative research. *J Adv Nurs.* 2021;77(9):3692-3706. doi:[10.1111/jan.14865](https://doi.org/10.1111/jan.14865)
25. Brooke J. SUS: A "Quick and Dirty" Usability Scale. In: *Usability Evaluation In Industry.* CRC Press; 1996.
26. Terwee CB, Prinsen CAC, Chiarotto A, et al. COSMIN methodology for evaluating the content validity of patient-reported outcome measures: a Delphi study. *Qual Life Res.* 2018;27(5):1159-1170. doi:[10.1007/s11136-018-1829-0](https://doi.org/10.1007/s11136-018-1829-0)
27. O'Brien HL, Cairns P, Hall M. A practical approach to measuring user engagement with the refined user engagement scale (UES) and new UES short form. *Int J Hum-Comput Stud.* 2018;112:28-39. doi:[10.1016/j.ijhcs.2018.01.004](https://doi.org/10.1016/j.ijhcs.2018.01.004)
28. Witmer BG, Singer MJ. Measuring presence in virtual environments: A presence questionnaire. *Presence Teleoperators Virtual Environ.* 1998;7(3):225-240. doi:[10.1162/105474698565686](https://doi.org/10.1162/105474698565686)
29. Spreng RN, McKinnon MC, Mar RA, Levine B. The Toronto Empathy Questionnaire: Scale Development and Initial Validation of a Factor-Analytic Solution to Multiple Empathy Measures. *J Pers Assess.* 2009;91(1):62-71. doi:[10.1080/00223890802484381](https://doi.org/10.1080/00223890802484381)
30. Norman G. Likert scales, levels of measurement and the "laws" of statistics. *Adv Health Sci Educ Theory Pract.* 2010;15(5):625-632. doi:[10.1007/s10459-010-9222-y](https://doi.org/10.1007/s10459-010-9222-y)
31. Wasserstein RL, Schirm AL, Lazar NA. Moving to a World Beyond "p < 0.05." *Am Stat.* 2019;73(sup1):1-19. doi:[10.1080/00031305.2019.1583913](https://doi.org/10.1080/00031305.2019.1583913)
32. Huk T, Ludwigs S. Combining cognitive and affective support in order to promote learning. *Learn Instr.* 2009;19(6):495-505. doi:[10.1016/j.learninstruc.2008.09.001](https://doi.org/10.1016/j.learninstruc.2008.09.001)
33. Park B, Plass JL, Brünken R. Cognitive and affective processes in multimedia learning. *Learn Instr.* 2014;29:125-127. doi:[10.1016/j.learninstruc.2013.05.005](https://doi.org/10.1016/j.learninstruc.2013.05.005)
34. Milota MM, Van Thiel GJMW, Van Delden JJM. Narrative medicine as a medical education tool: A systematic review. *Med Teach.* 2019;41(7):802-810. doi:[10.1080/0142159X.2019.1584274](https://doi.org/10.1080/0142159X.2019.1584274)
35. Lokka IE, Çöltekin A. Perspective switch and spatial knowledge acquisition: effects of age, mental rotation ability and visuospatial memory capacity on route learning in virtual environments with different levels of realism. *Cartogr Geogr Inf Sci.* 2020;47(1):14-27. doi:[10.1080/15230406.2019.1595151](https://doi.org/10.1080/15230406.2019.1595151)
36. Lokka IE, Çöltekin A, Wiener J, Fabrikant SI, Röcke C. Virtual environments as memory training devices in navigational tasks for older adults. *Sci Rep.* 2018;8(1):10809. doi:[10.1038/s41598-018-29029-x](https://doi.org/10.1038/s41598-018-29029-x)